



Oxidative modification of free-standing amino acids by Fe(II)/ α KG-dependent oxygenases

Hui Tao^{a,*}, Ikuro Abe^{a,b,*}

^a Graduate School of Pharmaceutical Sciences, The University of Tokyo, Tokyo, Japan

^b Collaborative Research Institute for Innovative Microbiology, The University of Tokyo, Tokyo, Japan

ARTICLE INFO

Keywords:

Fe(II)/ α KG-dependent oxygenases

Amino acids

Enzyme

ABSTRACT

Fe(II)/ α -ketoglutarate (α KG)-dependent oxygenases catalyze the oxidative modification of various molecules, from DNA, RNA, and proteins to primary and secondary metabolites. They also catalyze a variety of biochemical reactions, including hydroxylation, halogenation, desaturation, epoxidation, cyclization, peroxidation, epimerization, and rearrangement. Given the versatile catalytic capability of such oxygenases, numerous studies have been conducted to characterize their functions and elucidate their structure–function relationships over the past few decades. Amino acids, particularly nonproteinogenic amino acids, are considered as important building blocks for chemical synthesis and components for natural product biosynthesis. In addition, the Fe(II)/ α KG-dependent oxygenase superfamily includes important enzymes for generating amino acid derivatives, as they efficiently modify various free-standing amino acids. The recent discovery of new Fe(II)/ α KG-dependent oxygenases and the repurposing of known enzymes in this superfamily have promoted the generation of useful amino acid derivatives. Therefore, this study will focus on the recent progress achieved from 2019 to 2022 to provide a clear view of the mechanism by which these enzymes have expanded the repertoire of free amino acid oxidative modifications.

1. Introduction

Fe(II)- and α -ketoglutarate (α KG)-dependent oxygenases are widely distributed in nature, including microorganisms, plants, and vertebrates. During the past few decades, many studies have intensively investigated the mechanisms and structure–function relationships of Fe(II)/ α KG-dependent oxygenases. These efforts have revealed a generic mechanism and common double-stranded β -helix fold shared by the members of this superfamily [1,2,34,41,49,53,57,81,83]. Fe(II)/ α KG-dependent oxygenases catalyze the oxidative modification of primary and secondary metabolites, using O₂ with Fe(II) as the cofactor and α KG as the cosubstrate [40,43,59]. The oxidative decarboxylation of α KG to succinate generates a highly reactive Fe(IV)-oxo species, which selectively oxidizes inactivated C–H bonds and leads to a wide array of biochemical reactions, including hydroxylation, halogenation, desaturation, epoxidation, cyclization, peroxidation, epimerization, and rearrangement [10,17,22,25,33,56,63,65,68,71,85,92,97].

Given their association with numerous biological processes, Fe(II)/ α KG-dependent oxygenases play an important role in nature [36]. Many Fe(II)/ α KG-dependent oxygenases account for the modification of chromatin, DNA, and proteins, as well as the demethylation/hydroxylation of RNA [5,47,48,52,57]. Thus, such oxygenases have been reviewed recently [36]. Furthermore, given their versatile

catalytic roles and synthetic capabilities, Fe(II)/ α KG-dependent oxygenases can oxidize primary metabolites, such as amino acids, and tailor various biologically important natural products, including alkaloids and meroterpenoids [15,44,56,63,66,78,90,92]. The structure–function relationship of Fe(II)/ α KG-dependent oxygenases in the biosynthesis of natural products, particularly meroterpenoids, have been summarized in excellent reviews [7,26,28,36,54,64,84,91,98,104]. In addition, the characterization and engineering of halogenases have been summarized in reviews published in 2022 [74,96]. Thus, this study will focus on Fe(II)/ α KG-dependent oxygenases that are responsible for amino acid modifications.

Amino acid derivatives are considered as important nutritional supplements, pharmaceutical intermediates, and building blocks for organic synthesis and natural product biosynthesis [60,88,101]. Given their versatile applications, an increasing number of Fe(II)/ α KG-dependent oxygenases have been characterized for amino acid modifications. They oxidize carrier protein-bound and free-standing amino acids during natural product biosynthesis. For example, KtzO and KtzP cannot accept free-standing L-Glu-during kutzneride biosynthesis. Thus, they hydroxylate L-Glu-bound to a peptidyl carrier protein and generate L-threo- and L-erythro-hydroxyglutamic acids, respectively [87]. By contrast, some Fe(II)/ α KG-dependent hydroxylases accept free amino acids as substrates, such as the proline hydroxylase P3H/P4H [30,61,62], the

* Corresponding authors.

E-mail addresses: thuiwhu@gmail.com (H. Tao), abei@mol.f.u-tokyo.ac.jp (I. Abe).




Fig. 1. β -Hydroxylation of free-standing amino acids by (A) IboH, (B) GloE, and (C) AEP14369.

isoleucine 4-hydroxylase IDO [37,42,70,99], the leucine 5-hydroxylases LdoA and GriE [38,51,103], the lysine 3/4-hydroxylases KDO and K3H/K4H [8,32], the arginine 3-hydroxylase VioC [21,35,100], and the glutamine 3-hydroxylase AsnO [86]. These α KG-dependent amino acid hydroxylases have been described in previous reviews [26,39,53,74,75]. Therefore, we will primarily discuss the recent progress achieved from 2019 to 2022. Apart from summarizing the new Fe(II)/ α KG-dependent enzymes that were recently discovered and characterized for the oxidative modification of free-standing amino acids, we will also focus on reactions involving known repurposed enzymes, which facilitate the generation of useful amino acid derivatives.

2. Discovery of new Fe(II)/ α KG-dependent oxygenases to expand the repertoire of amino acid oxidative modifications

2.1. Hydroxylation of amino acids by Fe(II)/ α KG-dependent oxygenases

2.1.1. β -Hydroxylation

β -Hydroxy (β -OH or 3-OH) amino acids, including β -fluoro amino acids and β -lactams, are important precursors for pharmaceuticals and building blocks for organic synthesis [50,73]. In addition, β -OH amino acids are found in biologically active natural products, such as β -OH-Glu-in kutznerides [87] and β -OH-Arg-in viomycin [100]. Fe(II)/ α KG-dependent oxygenases are also important, as they not only modify amino acids tethered to a carrier protein, such as the generation of β -OH-Glu-bound to the peptidyl carrier protein by KtzO and KtzP during kutzneride biosynthesis [87], but also hydroxylate free-standing amino acids, such as L-Arg-by VioC [100] and L-Lys-by KDO1 [8]. However, until the characterization of IboH, which is the first reported Fe(II)/ α KG-dependent oxygenase for the hydroxylation of free L-Glu, no Fe(II)/ α KG-dependent oxygenase that could catalyze the β -hydroxylation of free-standing L-Glu had been identified [69] (Fig. 1A). IboH is encoded in the ibotenic acid biosynthetic gene cluster from *Amanita muscaria*. It accepts free L-Glu-to generate the product L-threo-3-OH-Glu, which ini-

tiates the biosynthesis of ibotenic acid in fly agaric. Moreover, IboH has a strict substrate preference, as L-Gln, which has a similar structure to L-Glu, was not accepted [69].

β -OH-Gln was discovered in the nonribosomal peptide pneumocandin B₀ [46]. In the biosynthetic gene cluster of pneumocandin B₀, an Fe(II)/ α KG-dependent oxygenase, GloE, was characterized as the β -hydroxylase that generates L-threo- β -OH-Gln from free-standing L-Gln (Fig. 1B) [79]. In addition to GloE, AEP14369 from *Sulfobacillus thermotolerans* Y0017 accepted L-Gln [31]. AEP14369 was characterized as an Fe(II)/ α KG-dependent β -hydroxylase that selectively generates L-threo- β -OH-Gln and L-threo- β -OH-His from L-Gln and L-His, respectively (Fig. 1C). β -OH-L-His is present in many natural products, including nikkomycin [18], siderophores [13,29], and bleomycin [89,95]. During nikkomycin biosynthesis, the heme- and NADPH-dependent protein NikQ catalyzes the β -hydroxylation of L-His-bound to a carrier protein domain [18]. The generation of β -OH-His in siderophores [77] and glidomides [19] is performed by Fe(II)/ α KG-dependent β -hydroxylases that use thiolation domain-bound L-His as a substrate. Notably, no Fe(II)/ α KG-dependent enzyme that accepts free-standing L-Gln and L-His had been reported until the discovery of GloE and AEP14369, which expanded the reaction repertoire of Fe(II)/ α KG-dependent β -hydroxylases.

2.1.2. γ -Hydroxylation

γ -Hydroxy (γ -OH or 4-OH) amino acids are biologically and pharmacologically important, as they are widely used as nutritional supplements (e.g., trans-4-OH-L-Pro), antidiabetic drugs (e.g., 4-OH-L-Ile, 4-OH-L-norVal, and 4-OH-L-Pip), and chiral building blocks for organic synthesis [60,101]. Furthermore, γ -OH-amino acids are important building blocks for natural products. Newly discovered Fe(II)/ α KG-dependent γ -hydroxylases have been characterized for the generation of 4-OH-L-Lys-in glidobactin [3], 4-OH-L-Gln-in gramillan A [79], and 4-OH-L-Cit in GE81112A [105] (Fig. 2).




Fig. 2. γ -Hydroxylation of free-standing amino acids by (A) GlbB, (B) Q4Ox, and (C) GetI.

Based on previous reports, L-Lys-hydroxylases such as KDO2, KDO3, and K4H produce (4R)-4-OH-L-Lys [8,32]. In contrast to these enzymes, an Fe(II)/ α KG-dependent oxygenase, GlbB, was found to selectively generate (4S)-4-OH-L-Lys during glidobactin biosynthesis (Fig. 2A) [3]. Biochemical characterization revealed that GlbB not only catalyzes the efficient γ -hydroxylation of free-standing L-Lys with complete regio- and diastereoselectivity, but also hydroxylates aliphatic amino acids (L-Leu- and L-Met) with moderate total turnover numbers [3]. Notably, Fe(II)/ α KG-dependent hydroxylases selectively act on aliphatic amino acids (e.g., IDO) [37,42,70] or polar amino acids (e.g., VioC) [21,35,100]. Therefore, GlbB is the first enzyme in this family that accepts both aliphatic and polar amino acids. Furthermore, the chemoenzymatic synthesis of cepafungin I, which is a potent inhibitor of the 20S proteasome core, was achieved through the 4-hydroxylation of L-Lys with GlbB [3].

4-OH-L-Gln was detected in nonribosomal peptide gramillan A, whose biosynthetic gene cluster contains the gene encoding the Fe(II)/ α KG-dependent oxygenase Q4Ox [79]. Biochemical characterizations revealed that Q4Ox produced (4R)-4-OH-L-Gln from free-standing L-Gln with complete diastereoselectivity (Fig. 2B), thereby providing a new enzyme for the preparation of useful glutamine derivatives.

GE81112s, which inhibit prokaryotic translation initiation, are nonribosomal tetrapeptides that contain several unusual amino acid building blocks, including 3-OH-L-Pip, 4-OH-L-Cit, and β -OH-2-Cl-L-His [12,24]. Two Fe(II)/ α KG-dependent oxygenases, GetI and GetF, were identified in the biosynthetic gene cluster of GE81112s [9]. In previous studies, GetF was characterized as an L-Pip β -hydroxylase, and GetI was annotated as a 2-Cl-L-His β -hydroxylase [9,55]. However, a recent study has revealed that GetI accepts L-Cit rather than 2-Cl-L-His to generate 4-OH-L-Cit [105] (Fig. 2C). Based on a homology model and sequence alignment analysis, the rational mutagenesis of GetI generated a vari-

ant that contains four mutations, as compared with a wild-type enzyme. Moreover, the GetI variant converted substrate preference from L-Cit to L-Arg. With the characterization of GetF and GetI, the chemoenzymatic total synthesis of GE81112 B1 and its analogs was completed, which promoted the first structure-activity studies of the antimicrobial activity of GE81112 [105].

2.2. Cyclization reactions by Fe(II)/ α KG-dependent oxygenases using free amino acids

2.2.1. Aziridination by Tqal

Several Fe(II)/ α KG-dependent oxygenases that accept free amino acids have been reported; however, most of them such as IboH [69], GloE [3], and AEP14369 [31] perform common hydroxylation reactions. Cyclization reactions catalyzed by enzymes in this superfamily, including CAS, H6H, HygX, LolO, and FfnD [22,25,33,56,71], have been reported. However, no enzyme was found to catalyze the cyclization of single and free-standing amino acids until the discovery of Tqal, which cyclizes L-Val to an aziridine-containing product during the biosynthesis of 2-aminoisobutyric acid (AIB), an important building block for bioactive natural products [14] (Fig. 3A). In the biosynthetic gene cluster of tryptoquialanine [27], Tqal was identified and characterized as an Fe(II)/ α KG-dependent oxygenase [14]. Tqal could initiate the reaction by abstracting the H-3 of L-Val via Fe(IV)-oxo species, leading to the generation of a C3-radical species. Subsequently, the C-N bond was formed to complete the aziridine-ring closure by either radical transfer or cation generation [14]. In addition, when the deuterated substrate (2S,3S)-[4,4,4-²H₃]-Val with a stereogenic center at C3 was used as a mechanistic probe, Tqal generated a diastereomeric pair of aziridines with the retention and inversion of C3 stereoconfiguration [93]. Furthermore, investigations on the stereoselectivity and substrate speci-




Fig. 3. Cyclization of free-standing amino acids to generate (A) aziridine ring from L-Val, (B) Ile-by TqaL, and (C) cyclopropane ring by BelL and HrmJ.

ficity of TqaL showed that nonnatural substrates, such as L-Ile and L-allo-Ile, were accepted by TqaL via a stereoconvergent process to generate diastereomeric pairs of aziridine and 3-OH-Ile-products (Fig. 3B), which is consistent with the mixed stereochemical course determined using (2S,3S)-[4,4,4-²H₃]-Val. Based on mutagenesis studies, the reaction type (aziridination *versus* hydroxylation) and stereochemical outcome of TqaL were regulated by two active site residues (I343 and F345). Furthermore, large-to-small substitution of F345 expanded the substrate scope of TqaL. For example, the F345S variant is highly active toward L-Leu- and L-Met- to generate hydroxy-L-Leu- and L-Met-sulfoxide. Although the detailed mechanism of the aziridine-ring closure requires

further investigation, these results have greatly expanded the catalytic repertoire of Fe(II)/ α KG-dependent oxygenases, thereby indicating that the stereochemical outcomes of the reactions could be rationally controlled by structure-guided mutagenesis.

2.2.2. Cyclopropanation by BelL and HrmJ

Natural products with strained three-membered rings, including aziridine and cyclopropane structures, such as aziridine-containing mitomycin and cyclopropane-containing belactosins and hormaomycins, have attracted great interest because of their potent biological activities [4,6,11,94]. In the structure of belactosins and hormaomycins, cyclo-




Fig. 4. Repurposing of (A) PolL and LdoA for nitrile or epoxide formation and (B) SmP4H for L-homophenylalanine oxidation. (C) The conversion of L-Lys-hydroxylase to Chi-14 halogenase.

propane rings were found in 3-(2-aminocyclopropyl)alanine (Acpa) and 3-(2-nitrocyclopropyl)alanine (Ncpa) residues, respectively [80,102]. Remarkably, Acpa and Ncpa exhibit opposite stereoconfigurations at C1' and C2' in the cyclopropane rings; thus, their biosynthesis has attracted great attention [23,45,72,82]. Two heme oxygenase-like diiron enzymes (BelK and HrmI, 51% identity) and two Fe(II)/ α KG-dependent oxygenases (BelL and HrmJ, 49% identity) were discovered by compar-

ing the biosynthetic gene clusters of belactosins and hormaomycins. In addition, *in vitro* and *in vivo* experiments revealed that BelK and HrmI oxidize L-Lys to 6-nitro-norLeu, which is subsequently accepted by BelL or HrmJ to generate Ncpa with opposite stereoconfigurations at the cyclopropane rings [45,72,82] (Fig. 3C). HrmJ produced (1'R,2'R)-Ncpa with high stereoselectivity, whereas BelL generated (1'S,2'S)-Ncp as the major product, which is consistent with the stereochemistry of cyclo-

Table 1
Substrate specificity and quantitative data of summarized enzymes.

	Substrates	Kinetics	TTNs	Refs.
IboH	L-Glu	–	–	[69]
GloE	L-Gln	–	44	[79]
AEP14369	L-Gln	$k_{cat} = 43.8 \pm 4.1 \text{ min}^{-1}$ $K_M = 1.17 \pm 0.27 \text{ mM}$	–	[31]
	L-His	$k_{cat} = 35.1 \pm 3.9 \text{ min}^{-1}$ $K_M = 0.83 \pm 0.06 \text{ mM}$	–	
GlbB	L-Lys	$k_{cat} = 960 \pm 84 \text{ min}^{-1}$ $K_M = 0.034 \pm 0.005 \text{ mM}$	5900	[3]
	L-Leu	$k_{cat} = 3.2 \pm 0.2 \text{ min}^{-1}$ $K_M = 2.2 \pm 0.4 \text{ mM}$	310	
	L-Met	$k_{cat} = 12.0 \pm 0.3 \text{ min}^{-1}$ $K_M = 0.45 \pm 0.05 \text{ mM}$	330	
Q4Ox	L-Gln	$k_{cat} = 250 \pm 3 \text{ min}^{-1}$ $K_M = 0.99 \pm 0.03 \text{ mM}$	4300	[79]
Getf	L-Cit	$k_{cat} = 69.0 \pm 5.3 \text{ min}^{-1}$ $K_M = 2.1 \pm 0.4 \text{ mM}$	470	[105]
	L-Arg	$k_{cat} = 250 \pm 3 \text{ min}^{-1}$ $K_M = 0.99 \pm 0.03 \text{ mM}$	20	
Tqal	L-Val	$k_{cat} = 22.1 \pm 1.0 \text{ min}^{-1}$ $K_M = 172 \pm 25 \mu\text{M}$	–	[93]
	L-Ile	$k_{cat} = 14.5 \pm 0.4 \text{ min}^{-1}$ $K_M = 665 \pm 45 \mu\text{M}$	–	
	L- <i>allo</i> -Ile	$k_{cat} = 0.32 \pm 0.03 \text{ min}^{-1}$ $K_M = 270 \pm 73 \mu\text{M}$	–	
HrmJ	6-nitronorleucine	–	–	[82]
BelL	6-nitronorleucine	–	–	[45]
PolL	(2S)-2-amino-4-azidobutanoic acid	$k_{cat} = 4.08 \pm 0.74 \text{ min}^{-1}$ $K_M = 1.49 \pm 0.65 \text{ mM}$	150	[20]
LdoA	5-azido-L-norvaline	$k_{cat} = 1.90 \pm 0.18 \text{ min}^{-1}$ $K_M = 1.30 \pm 0.32 \text{ mM}$	180	
	(S)-2-aminohex-5-enoic acid	–	–	[16]
Smp4H (WT)	L-hPhe	$k_{cat} = 0.015 \pm 0.001 \text{ min}^{-1}$ $K_M = 1.10 \pm 0.24 \text{ mM}$	4	[58]
Smp4H/ W40M I103L	L-hPhe	$k_{cat} = 1.680 \pm 0.068 \text{ min}^{-1}$ $K_M = 0.40 \pm 0.08 \text{ mM}$	39	
Hydrox	L-Lys	$k_{cat} = 22.5 \pm 0.5 \text{ min}^{-1}$ $K_M = 0.34 \pm 0.05 \text{ mM}$	136	[74]
Hal	L-Lys	$k_{cat} = 10.5 \pm 0.4 \text{ min}^{-1}$ $K_M = 0.12 \pm 0.03 \text{ mM}$	410	
Chi-14	L-Lys	$k_{cat} = 20.3 \pm 0.5 \text{ min}^{-1}$ $K_M = 0.29 \pm 0.03 \text{ mM}$	410	

Note: "–" indicates no data has been reported yet. TTNs: Total turnover numbers.

propane rings found in hormaomycins and belactosins. Investigations of the H abstraction site, which used stereoselectively deuterated 6-nitronorLeu, revealed that BelL and HrmJ selectively abstract 4-*pro*S-H, but they exhibited no or little stereoselectivity for the dehydrogenation at C6 [82]. Although additional investigations are necessary to understand the mechanism by which these enzymes control the stereochemistry, these observations have led to the identification of a novel approach for the stereoselective construction of cyclopropane rings in natural products.

3. Repurposing Fe(II)/ α KG-dependent oxygenases for the synthesis of new amino acid derivatives

The discovery and characterization of new members in the Fe(II)/ α KG-dependent oxygenase superfamily significantly expanded the catalytic repertoire. Recently, the repurposing of previously characterized enzymes, such as LdoA, PolL, and Smp4H, has emerged as a novel approach for generating new amino acid derivatives [16,20,58] (Fig. 4). LdoA from *Nostoc punctiforme* ACC80786 and PolL from *Streptomyces aureochromogenes* are considered as Fe(II)/ α KG-dependent oxygenases that catalyze the hydroxylation of L-Leu-at the C5 position [38] and the dihydroxylation of L- α -amino- δ -carbamoylhydroxyvaleric acid at the C3 and C4 positions [76], respectively. In a recent study, LdoA and PolL were found to catalyze nitrile formation when an azido group was introduced into nonnative substrates [20] (Fig. 4A). Moreover, when an olefin group was installed on the LdoA substrate, the reaction was redirected to asymmetric epoxidation, which led to the generation of 5-OH-

Pip [16] (Fig. 4A). Smp4H, a proline 4-hydroxylase from *Sinorhizobium meliloti*, was repurposed as an L-homophenylalanine (hPhe) hydroxylase [58] (Fig. 4B). Furthermore, structure-based rational engineering identified the W40M I103L variant with increased activity to generate further oxidized products, including 4-OH-hPhe, 4-ketone-hPhe, and 3-OH-4-ketone-hPhe (Fig. 4B, Table 1). Notably, with only a single-site mutation, the W40Y variant primarily served as a desaturase [58] (Fig. 4B). These studies revealed the catalytic potential of Fe(II)/ α KG-dependent oxygenases in synthetic and industrial applications.

Compared with the wide range of Fe(II)/ α KG-dependent hydroxylases, the number of halogenases in this superfamily is limited. Recently, the engineering of an Fe(II)/ α KG-dependent lysine hydroxylase into a halogenase was achieved [67] (Fig. 4C). Based on the phylogenetic tree of the BesD family, hydroxylases and halogenases are grouped into separate clades. However, a hydroxylase (Hydrox) was included in the clade of putative halogenases. The Hydrox enzyme shares 71% amino acid sequence identity with a halogenase (Hal). Biochemical characterization revealed that Hydrox and Hal catalyze the hydroxylation and chlorination of L-Lys-at the C4 position, respectively. Hydroxylase-halogenase chimeric enzymes were generated by constructing a DNA shuffling library using these two genes, and halogenation-active variants were screened using a high-throughput *in vivo* fluorescent screen strategy. Finally, a variant (Chi-14) with 14 mutations was identified with comparable activity and higher selectivity to the wild-type halogenase Hal (Fig. 4C and Table 1). Most of the key mutations are located on the two β -sheets lining the α KG binding pocket, except for the criti-

cal residue Gly144 that enables chloride coordination to the Fe(IV)-oxo species, thereby indicating the importance of second-sphere residues for protein engineering, as they further tuned the activity and selectivity of the engineered enzyme [67]. This study promotes the conversion of hydroxylases to halogenases, thereby leading to the expansion of enzymatic halogenation.

4. Conclusions

Recently, an increasing number of Fe(II)/ α KG-dependent oxygenases were identified and well-characterized, and reaction repurposing was presented as a useful approach for the generation of important amino acid derivatives. The characterization of new Fe(II)/ α KG-dependent oxygenases enriched the toolbox for amino acid hydroxylation and cyclization. However, a number of putative Fe(II)/ α KG-dependent oxygenases remain unknown, which need to be characterized. Repurposing the function of known Fe(II)/ α KG-dependent oxygenases could be achieved by protein engineering and substrate design. For these newly discovered enzymes, although their functions and substrate scopes were characterized, the detailed mechanism and structure–function relationship need further investigations. This could significantly facilitate the rational protein engineering and design of nonnatural substrates. Furthermore, high-throughput screening and machine learning would assist enzyme engineering and boost the development of Fe(II)/ α KG-dependent oxygenases as potential catalysts for chemosynthetic and industrial application.

Declaration of Competing Interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Given his role as Editorial Board Member, Dr. Ikuro Abe, had no involvement in the peer-review of this article and has no access to information regarding its peer-review. Full responsibility for the editorial process for this article was delegated to Dr. Wei Zhang.

Acknowledgments

The authors would like to express sincere appreciation to an excellent group of coworkers whose contributions are cited in the text. We thank Drs. Takayoshi Awakawa, Takahiro Mori, and Richiro Ushimaru for critical reading of the paper. This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan (JSPS KAKENHI Grant No. JP16H06443, JP20KK013, and JP20H00490), and Japan Science and Technology Agency (JST SICORP Grant No. JPMJSC1701), the New Energy and Industrial Technology Development Organization (NEDO, Grant No. JPNP20011), and Japan Agency for Medical Research and Development (AMED) (Grant No. JP21ak0101164). H.T. is a recipient of the JSPS Postdoctoral Fellowship for Foreign Researchers (ID No. P18404).

References

- [1] W. Aik, M.A. McDonough, A. Thalhammer, et al., Role of the jelly-roll fold in substrate binding by 2-oxoglutarate oxygenases, *Curr. Opin. Struct. Biol.* 22 (6) (2012) 691–700.
- [2] W.S. Aik, R. Chowdhury, I.J. Clifton, et al., Introduction to structural studies on 2-oxoglutarate-dependent oxygenases and related enzymes. 2-Oxoglutarate-dependent oxygenases, *Z. Naturforsch., C, J. Biosci.* 45 (7–8) (1990) 851–855.
- [3] A. Amatuni, H. Renata, Identification of a lysine 4-hydroxylase from the glidobactin biosynthesis and evaluation of its biocatalytic potential, *Org. Biomol. Chem.* 17 (7) (2019) 1736–1739.
- [4] N. Andres, H. Wolf, H. Zähler, Hormaomycin, a new peptide lactone antibiotic effective in inducing cytodifferentiation and antibiotic biosynthesis in some *Streptomyces* species, *Z. Naturforsch., C, J. Biosci.* 45 (7–8) (1990) 851–855.
- [5] L. Aravind, E.V. Koonin, The DNA-repair protein AlkB, EGL-9, and Iprecan define new families of 2-oxoglutarate- and iron-dependent dioxygenases, *Genome Biol.* 2 (3) (2001) 1–8.

- [6] A. Asai, A. Hasegawa, K. Ochiai, et al., Belactosin A, a novel antitumor antibiotic acting on cyclin/CDK mediated cell cycle regulation, produced by *Streptomyces* sp, *J. Antibiot.* 53 (1) (2000) 81–83.
- [7] T. Awakawa, T. Mori, R. Ushimaru, et al., Structure-based engineering of α -ketoglutarate dependent oxygenases in fungal meroterpenoid biosynthesis, *Nat. Prod. Rep.* (2022).
- [8] D. Baud, P.-L. Saaidi, A. Monfleur, et al., Synthesis of mono- and dihydroxylated amino acids with new α -ketoglutarate-dependent dioxygenases: biocatalytic oxidation of C-H bonds, *ChemCatChem* 6 (10) (2014) 3012–3017.
- [9] T.M. Binz, S.I. Maffioli, M. Sosio, et al., Insights into an unusual nonribosomal peptide synthetase biosynthesis: identification and characterization of the GE81112 biosynthetic gene cluster, *J. Biol. Chem.* 285 (43) (2010) 32710–32719.
- [10] L.C. Blasiak, F.H. Vaillancourt, C.T. Walsh, et al., Crystal structure of the non-haem iron halogenase SyrB2 in syringomycin biosynthesis, *Nature* 440 (7082) (2006) 368–371.
- [11] W.T. Bradner, Mitomycin C: a clinical update, *Cancer Treat. Rev.* 27 (1) (2001) 35–50.
- [12] L. Brandi, A. Fabbretti, A. La Teana, et al., Specific, efficient, and selective inhibition of prokaryotic translation initiation by a novel peptide antibiotic, *Proc. Natl. Acad. Sci. U.S.A.* 103 (1) (2006) 39–44.
- [13] H. Budzikiewicz, S. Kilz, K. Taraz, et al., Identical Pyoverdines from *Pseudomonas fluorescens* 9AW and from *Pseudomonas putida* 9BW. *Z. Naturforsch., C, J. Biosci.* 52 (11–12) (1997) 721–728.
- [14] R. Bunno, T. Awakawa, T. Mori, et al., Aziridine formation by a Fe(II)/ α -ketoglutarate dependent oxygenase and 2-aminoisobutyrate biosynthesis in fungi, *Angew. Chem. Int. Ed.* 60 (29) (2021) 15827–15831.
- [15] P.D. Cárdenas, P.D. Sonawane, U. Heinig, et al., Pathways to defense metabolites and evading fruit bitterness in genus *Solanum* evolved through 2-oxoglutarate-dependent dioxygenases, *Nat. Commun.* 10 (1) (2019) 1–13.
- [16] L. Cha, S. Milikisiyants, M. Davidson, et al., Alternative reactivity of leucine 5-hydroxylase using an olefin-containing substrate to construct a substituted piperidine ring, *Biochemistry* 59 (21) (2020) 1961–1965.
- [17] W.-C. Chang, Y. Guo, C. Wang, et al., Mechanism of the C5 stereoinversion reaction in the biosynthesis of carbapenem antibiotics, *Science* 343 (6175) (2014) 1140–1144.
- [18] H. Chen, B.K. Hubbard, S.E. O'Connor, et al., Formation of β -hydroxy histidine in the biosynthesis of nikkomycin antibiotics, *Chem. Biol.* 9 (1) (2002) 103–112.
- [19] H. Chen, L. Zhong, H. Zhou, et al., Biosynthesis of glidomides and elucidation of different mechanisms for formation of β -OH amino acid building blocks, *Angew. Chem. Int. Ed.* 134 (35) (2022) e202203591.
- [20] M. Davidson, M. McNamee, R. Fan, et al., Repurposing nonheme iron hydroxylases to enable catalytic nitrile installation through an azido group assistance, *J. Am. Chem. Soc.* 141 (8) (2019) 3419–3423.
- [21] N.P. Dunham, A.J. Mitchell, J.M. Del Río Pantoja, et al., α -Amine desaturation of D-arginine by the iron(II)- and 2-(oxo)glutarate-dependent L-arginine 3-hydroxylase, *VioC, Biochemistry* 57 (46) (2018) 6479–6488.
- [22] S.W. Elson, K.H. Baggaley, J. Gillett, et al., Isolation of two novel intracellular β -lactams and a novel dioxygenase cyclising enzyme from *Streptomyces clavuligerus*, *J. Chem. Soc. Chem. Commun.* 22 (1987) 1736–1738.
- [23] A. Engelbrecht, F. Wolf, A. Esch, et al., Discovery of a cryptic nitro intermediate in the biosynthesis of the 3-(trans-2'-aminocyclopropyl)alanine moiety of belactosin A, *Org. Lett.* 24 (2) (2022) 736–740.
- [24] A. Fabbretti, A. Schedlbauer, L. Brandi, et al., Inhibition of translation initiation complex formation by GE81112 unravels a 16S rRNA structural switch involved in P-site decoding, *Proc. Natl. Acad. Sci. U.S.A.* 113 (16) (2016) E2286–E2295.
- [25] S.R. Gama, T. Stankovic, K. Hupp, et al., Biosynthesis of the fungal organophosphate fosfonochlorin involves an iron (II) and 2-(oxo) glutarate dependent oxacyclase, *ChemBioChem* 23 (2) (2022) e202100352.
- [26] S.-S. Gao, N. Naowarajna, R. Cheng, et al., Recent examples of α -ketoglutarate-dependent mononuclear non-haem iron enzymes in natural product biosynthesis, *Nat. Prod. Rep.* 35 (8) (2018) 792–837.
- [27] X. Gao, Y.-H. Chooi, B.D. Ames, et al., Fungal indole alkaloid biosynthesis: genetic and biochemical investigation of the tryptotoqualanine pathway in *Penicillium aethiopicum*, *J. Am. Chem. Soc.* 133 (8) (2011) 2729–2741.
- [28] R. Geris, T.J. Simpson, Meroterpenoids produced by fungi, *Nat. Prod. Rep.* 26 (8) (2009) 1063–1094.
- [29] D.K. Hancock, B. Coxon, S.-Y. Wang, et al., L-threo- β -hydroxyhistidine, an unprecedented iron(III) ion-binding amino acid in a pyoverdine-type siderophore from *Pseudomonas fluorescens* 244, *J. Chem. Soc. Chem. Commun.* (5) (1993) 468–470.
- [30] R. Hara, K. Kino, Characterization of novel 2-oxoglutarate dependent dioxygenases converting L-proline to cis-4-hydroxy-L-proline, *Biochem. Biophys. Res. Commun.* 379 (4) (2009) 882–886.
- [31] R. Hara, Y. Nakajima, H. Yanagawa, et al., Enzymatic synthesis of L-threo- β -hydroxy- α -amino acids via asymmetric hydroxylation using 2-oxoglutarate-dependent hydroxylase from *Sulfobacillus thermotolerans* strain Y0017, *Appl. Environ. Microbiol.* 87 (20) (2021) e01335–01321.
- [32] R. Hara, K. Yamagata, R. Miyake, et al., Discovery of lysine hydroxylases in the clavaminic acid synthase-like superfamily for efficient hydroxylysine bioproduction, *Appl. Environ. Microbiol.* 83 (17) (2017) e00693–00617.
- [33] T. Hashimoto, A. Hayashi, Y. Amano, et al., Hyoscyamine 6 β -hydroxylase, an enzyme involved in tropane alkaloid biosynthesis, is localized at the pericycle of the root, *J. Biol. Chem.* 266 (7) (1991) 4648–4653.
- [34] Hausinger, R.P. 2015. Biochemical diversity of 2-oxoglutarate-dependent oxygenases.

- [35] V. Helmetag, S.A. Samel, M.G. Thomas, et al., Structural basis for the erythro-stereospecificity of the L-arginine oxygenase VioC in viomycin biosynthesis, *FEBS J.* 276 (13) (2009) 3669–3682.
- [36] C.Q. Herr, R.P. Hausinger, Amazing diversity in biochemical roles of Fe (II)/2-oxoglutarate oxygenases, *Trends Biochem. Sci.* 43 (7) (2018) 517–532.
- [37] M. Hibi, T. Kawashima, T. Kodera, et al., Characterization of *Bacillus thuringiensis* L-isoleucine dioxygenase for production of useful amino acids, *Appl. Environ. Microbiol.* 77 (19) (2011) 6926–6930.
- [38] M. Hibi, T. Kawashima, P.M. Sokolov, et al., L-Leucine 5-hydroxylase of *Nostoc punctiforme* is a novel type of Fe (II)/ α -ketoglutarate-dependent dioxygenase that is useful as a biocatalyst, *Appl. Microbiol. Biotechnol.* 97 (6) (2013) 2467–2472.
- [39] M. Hibi, J. Ogawa, Characteristics and biotechnology applications of aliphatic amino acid hydroxylases belonging to the Fe(II)/ α -ketoglutarate-dependent dioxygenase superfamily, *Appl. Microbiol. Biotechnol.* 98 (9) (2014) 3869–3876.
- [40] X. Huang, J.T. Groves, Beyond ferryl-mediated hydroxylation: 40 years of the rebound mechanism and C-H activation, *J. Biol. Inorg. Chem.* 22 (2) (2017) 185–207.
- [41] M.S. Islam, T.M. Leissing, R. Chowdhury, et al., 2-Oxoglutarate-dependent oxygenases, *Annu. Rev. Biochem.* 87 (1) (2018) 585–620.
- [42] T. Kodera, S.V. Smirnov, N.N. Samsonova, et al., A novel L-isoleucine hydroxylating enzyme, L-isoleucine dioxygenase from *Bacillus thuringiensis*, produces (2S, 3R, 4S)-4-hydroxyisoleucine, *Biochem. Biophys. Res. Commun.* 390 (3) (2009) 506–510.
- [43] C. Krebs, D. Galonic Fujimori, C.T. Walsh, et al., Non-heme Fe (IV)-oxo intermediates, *Acc. Chem. Res.* 40 (7) (2007) 484–492.
- [44] X. Li, T. Awakawa, T. Mori, et al., Heterodimeric non-heme iron enzymes in fungal meroterpenoid biosynthesis, *J. Am. Chem. Soc.* 143 (50) (2021) 21425–21432.
- [45] X. Li, R. Shimaya, T. Dairi, et al., Identification of cyclopropane formation in the biosyntheses of hormaomycins and belactosins: sequential nitration and cyclopropanation by metalloenzymes, *Angew. Chem. Int. Ed.* 61 (7) (2022) e202113189.
- [46] Y. Li, N. Lan, L. Xu, et al., Biosynthesis of pneumocandin lipopeptides and perspectives for its production and related echinocandins, *Appl. Microbiol. Biotechnol.* 102 (23) (2018) 9881–9891.
- [47] C. Loenarz, C.J. Schofield, Oxygenase catalyzed 5-methylcytosine hydroxylation, *Chem. Biol.* 16 (6) (2009) 580–583.
- [48] C. Loenarz, C.J. Schofield, Physiological and biochemical aspects of hydroxylations and demethylations catalyzed by human 2-oxoglutarate oxygenases, *Trends Biochem. Sci.* 36 (1) (2011) 7–18.
- [49] J.E. Longbotham, C. Levy, L.O. Johannissen, et al., Structure and mechanism of a viral collagen prolyl hydroxylase, *Biochemistry* 54 (39) (2015) 6093–6105.
- [50] B.T. Lotz, M.J. Miller, Diastereoselective synthesis of the carbacephem framework, *J. Org. Chem.* 58 (3) (1993) 618–625.
- [51] P. Lukat, Y. Katsuyama, S. Wenzel, et al., Biosynthesis of methyl-proline containing griselimycins, natural products with anti-tuberculosis activity, *Chem. Sci.* 8 (11) (2017) 7521–7527.
- [52] S. Markolovic, T.M. Leissing, R. Chowdhury, et al., Structure–function relationships of human JmjC oxygenases—Demethylases versus hydroxylases, *Curr. Opin. Struct. Biol.* 41 (2016) 62–72.
- [53] S. Martinez, R.P. Hausinger, Catalytic mechanisms of Fe (II)- and 2-oxoglutarate-dependent oxygenases, *J. Biol. Chem.* 290 (34) (2015) 20702–20711.
- [54] Y. Matsuda, I. Abe, 1,14-Fungal meroterpenoids, in: *Comprehensive Natural Products II*, 1, Elsevier, Oxford, 2020, pp. 445–478.
- [55] J. Mattay, W. Hüttel, Pipecolic acid hydroxylases: a monophyletic clade among cis-selective bacterial proline hydroxylases that discriminates L-proline, *ChemBioChem* 18 (15) (2017) 1523–1528.
- [56] K.M. McCulloch, E.K. McCranie, J.A. Smith, et al., Oxidative cyclizations in orthosomycin biosynthesis expand the known chemistry of an oxygenase superfamily, *Proc. Natl. Acad. Sci. U.S.A.* 112 (37) (2015) 11547–11552.
- [57] M.A. McDonough, C. Loenarz, R. Chowdhury, et al., Structural studies on human 2-oxoglutarate dependent oxygenases, *Curr. Opin. Struct. Biol.* 20 (6) (2010) 659–672.
- [58] F. Meyer, R. Frey, M. Ligibel, et al., Modulating chemoselectivity in a Fe(II)/ α -ketoglutarate-dependent dioxygenase for the oxidative modification of a nonproteinogenic amino acid, *ACS Catal.* 11 (10) (2021) 6261–6269.
- [59] A.J. Mitchell, N.P. Dunham, R.J. Martinie, et al., Visualizing the reaction cycle in an iron(ii)- and 2-(oxo)-glutarate-dependent hydroxylase, *J. Am. Chem. Soc.* 139 (39) (2017) 13830–13836.
- [60] C.J. Moreno, K. Hernández, S.J. Charnok, et al., Synthesis of γ -hydroxy- α -amino acid derivatives by enzymatic tandem aldol addition–transamination reactions, *ACS Catal.* 11 (8) (2021) 4660–4669.
- [61] H. Mori, T. Shibasaki, Y. Uozaki, et al., Detection of novel proline 3-hydroxylase activities in *Streptomyces* and *Bacillus* spp. by regio- and stereospecific hydroxylation of L-proline, *Appl. Environ. Microbiol.* 62 (6) (1996) 1903–1907.
- [62] H. Mori, T. Shibasaki, K. Yano, et al., Purification and cloning of a proline 3-hydroxylase, a novel enzyme which hydroxylates free L-proline to cis-3-hydroxy-L-proline, *J. Bacteriol.* 179 (18) (1997) 5677–5683.
- [63] T. Mori, R. Zhai, R. Ushimaru, et al., Molecular insights into the endoperoxide formation by Fe (II)/ α -KG-dependent oxygenase Nvfl, *Nat. Commun.* 12 (1) (2021) 1–11.
- [64] H. Nakamura, Y. Matsuda, I. Abe, Unique chemistry of non-heme iron enzymes in fungal biosynthetic pathways, *Nat. Prod. Rep.* 35 (7) (2018) 633–645.
- [65] Y. Nakashima, T. Mori, H. Nakamura, et al., Structure function and engineering of multifunctional non-heme iron dependent oxygenases in fungal meroterpenoid biosynthesis, *Nat. Commun.* 9 (1) (2018) 1–10.
- [66] R.S. Nett, Y. Dho, Y.-Y. Low, et al., A metabolic regulon reveals early and late acting enzymes in neuroactive Lycopodium alkaloid biosynthesis, *Proc. Natl. Acad. Sci. U.S.A.* 118 (24) (2021) e2102949118.
- [67] M.E. Neugebauer, E.N. Kissman, J.A. Marchand, et al., Reaction pathway engineering converts a radical hydroxylase into a halogenase, *Nat. Chem. Biol.* 18 (2) (2022) 171–179.
- [68] M.E. Neugebauer, K.H. Sumida, J.G. Pelton, et al., A family of radical halogenases for the engineering of amino-acid-based products, *Nat. Chem. Biol.* 15 (10) (2019) 1009–1016.
- [69] S. Obermaier, M. Müller, Ibotenic acid biosynthesis in the fly agaric is initiated by glutamate hydroxylation, *Angew. Chem. Int. Ed.* 132 (30) (2020) 12532–12535.
- [70] J. Ogawa, T. Kodera, S.V. Smirnov, et al., A novel L-isoleucine metabolism in *Bacillus thuringiensis* generating (2S, 3R, 4S)-4-hydroxyisoleucine, a potential insulinotropic and anti-obesity amino acid, *Appl. Microbiol. Biotechnol.* 89 (6) (2011) 1929–1938.
- [71] J. Pan, M. Bhardwaj, B. Zhang, et al., Installation of the ether bridge of lolines by the iron- and 2-oxoglutarate-dependent oxygenase, LolO: regio- and stereochemistry of sequential hydroxylation and oxacyclization reactions, *Biochemistry* 57 (14) (2018) 2074–2083.
- [72] L. Pang, W. Niu, Y. Duan, et al., *In vitro* characterization of a nitro-forming oxygenase involved in 3-(trans-2'-aminocyclopropyl)alanine biosynthesis, *Eng. Microbiol.* 2 (1) (2022) 100007.
- [73] S.V. Pansare, J.C. Vederas, Reaction of β -hydroxy α -amino acid derivatives with (diethylamino) sulfur trifluoride (DAST). Synthesis of β -fluoro α -amino acids, *J. Org. Chem.* 52 (21) (1987) 4804–4810.
- [74] A. Papadopoulou, F. Meyer, R.M. Buller, Engineering Fe(II)/ α -ketoglutarate-dependent halogenases and desaturases, *Biochemistry* (2022).
- [75] C. Peters, R.M. Buller, Industrial application of 2-oxoglutarate-dependent oxygenases, *Catalysts* 9 (3) (2019) 221.
- [76] J. Qi, D. Wan, H. Ma, et al., Deciphering carbamoylpolyoamic acid biosynthesis reveals unusual acetylation cycle associated with tandem reduction and sequential hydroxylation, *Cell Chem. Biol.* 23 (8) (2016) 935–944.
- [77] Z.L. Reitz, C.D. Hardy, J. Suk, et al., Genomic analysis of siderophore β -hydroxylases reveals divergent stereocontrol and expands the condensation domain family, *Proc. Natl. Acad. Sci. U.S.A.* 116 (40) (2019) 19805–19814.
- [78] D. Ren, Y.-H. Lee, S.-A. Wang, et al., Characterization of the oxazinomycin biosynthetic pathway revealing the key role of a nonheme iron-dependent mono-oxygenase, *J. Am. Chem. Soc.* 144 (24) (2022) 10968–10977.
- [79] H. Renata, E. Shimizu, C.R. Zwick, Regiodivergent biocatalytic hydroxylation of L-glutamine facilitated by characterization of non-heme dioxygenases from non-ribosomal peptide biosyntheses, *Tetrahedron* 90 (2021) 132190.
- [80] E. Rössner, A. Zeek, W.A. König, Elucidation of the structure of hormaomycin, *Angew. Chem. Int. Ed.* 29 (1) (1990) 64–65.
- [81] M.J. Ryle, R.P. Hausinger, Non-heme iron oxygenases, *Curr. Opin. Chem. Biol.* 6 (2) (2002) 193–201.
- [82] S. Shimo, R. Ushimaru, A. Engelbrecht, et al., Stereodivergent nitrocyclopropane formation during biosynthesis of belactosins and hormaomycins, *J. Am. Chem. Soc.* 143 (44) (2021) 18413–18418.
- [83] T.J. Smart, R.B. Hamed, T.D.W. Claridge, et al., Studies on the selectivity of proline hydroxylases reveal new substrates including bicycles, *Bioorg. Chem.* 94 (2020) 103386.
- [84] H. Song, N. Naowarajna, R. Cheng, et al., Non-heme iron enzyme-catalyzed complex transformations: endoperoxidation, cyclopropanation, orthoester, oxidative CC and CS bond formation reactions in natural product biosynthesis, *Adv. Protein Chem. Struct. Biol.* 117 (2019) 1–61.
- [85] N. Steffan, A. Grundmann, S. Afiyatullo, et al., FtmOx1, a non-heme Fe (II) and α -ketoglutarate-dependent dioxygenase, catalyses the endoperoxide formation of verruculogen in *Aspergillus fumigatus*, *Org. Biomol. Chem.* 7 (19) (2009) 4082–4087.
- [86] M. Strieker, F. Kopp, C. Mahler, et al., Mechanistic and structural basis of stereospecific C β -hydroxylation in calcium-dependent antibiotic, a daptomycin-type lipopeptide, *ACS Chem. Biol.* 2 (3) (2007) 187–196.
- [87] M. Strieker, E.M. Nolan, C.T. Walsh, et al., Stereospecific synthesis of *threo*- and *erythro*- β -hydroxyglutamic acid during kutzneride biosynthesis, *J. Am. Chem. Soc.* 131 (37) (2009) 13523–13530.
- [88] P. Szczesniak, A. Pazzierniok-Holewa, U. Klimczak, et al., Synthesis of β - and γ -hydroxy α -amino acids via enzymatic kinetic resolution and cyanate-to-isocyanate rearrangement, *J. Org. Chem.* 79 (23) (2014) 11700–11713.
- [89] O. Takashi, O. Masami, O. Masaji, Enantioselective synthesis of *erythro*- β -Hydroxy-L-histidine, the pivotal amino acid of bleomycin-Fe(II)-O₂ complex, *Chem. Lett.* 17 (1) (1988) 83–86.
- [90] H. Tang, M.-H. Wu, H.-Y. Lin, et al., Mechanistic analysis of carbon-carbon bond formation by deoxypodophyllotoxin synthase, *Proc. Natl. Acad. Sci. U.S.A.* 119 (1) (2022) e2113770119.
- [91] H. Tao, I. Abe, Harnessing Fe (II)/ α -ketoglutarate-dependent oxygenases for structural diversification of fungal meroterpenoids, *Curr. Opin. Biotechnol.* 77 (2022) 102763.
- [92] H. Tao, T. Mori, H. Chen, et al., Molecular insights into the unusually promiscuous and catalytically versatile Fe (II)/ α -ketoglutarate-dependent oxygenase SptF, *Nat. Commun.* 13 (1) (2022) 1–13.
- [93] H. Tao, R. Ushimaru, T. Awakawa, et al., Stereoselectivity and substrate specificity of the Fe(II)/ α -ketoglutarate-dependent oxygenase Tqal, *J. Am. Chem. Soc.* 144 (2022) in press.
- [94] C.J. Thibodeaux, W.-C. Chang, H.-W. Liu, Enzymatic chemistry of cyclopropane, epoxide, and aziridine biosynthesis, *Chem. Rev.* 112 (3) (2012) 1681–1709.
- [95] M.G. Thomas, M.D. Burkart, C.T. Walsh, Conversion of L-proline to pyrrolyl-2-carboxyl-S-PCP during undecylprodigiosin and pyoluteorin biosynthesis, *Chem. Biol.* 9 (2) (2002) 171–184.

- [96] H.B. Wang, C. Che, S. You, Recent advances of enzymatic synthesis of organohalogen catalyzed by Fe/ α KG-dependent halogenases, *Synth. Biol. J.* 3 (3) (2022) 545.
- [97] S.D. Wong, M. Srnc, M.L. Matthews, et al., Elucidation of the Fe (IV)=O intermediate in the catalytic cycle of the halogenase SyrB2, *Nature* 499 (7458) (2013) 320–323.
- [98] L.-F. Wu, S. Meng, G.-L. Tang, Ferrous iron and α -ketoglutarate-dependent dioxygenases in the biosynthesis of microbial natural products, *Biochim. Biophys. Acta - Proteins Proteom.* 1864 (5) (2016) 453–470.
- [99] L. Wu, J. An, X. Jing, et al., Molecular insights into the regioselectivity of the Fe (II)/2-Ketoglutarate-dependent dioxygenase-catalyzed C–H hydroxylation of amino acids, *ACS Catal.* 12 (2022) 11586–11596.
- [100] X. Yin, T.M. Zabriske, VioC is a non-heme iron, α -ketoglutarate-dependent oxygenase that catalyzes the formation of 3S-hydroxy-L-arginine during viomycin biosynthesis, *ChemBioChem* 5 (9) (2004) 1274–1277.
- [101] M.I. Zafar, F. Gao, 4-Hydroxyisoleucine: a potential new treatment for type 2 diabetes mellitus, *Biodrugs* 30 (4) (2016) 255–262.
- [102] B.D. Zlatopolskiy, K. Loscha, P. Alvermann, et al., Final elucidation of the absolute configuration of the signal metabolite hormaomycin, *Eur. J. Chem.* 10 (19) (2004) 4708–4717.
- [103] C.R. Zwick, H. Renata, Remote C–H hydroxylation by an α -ketoglutarate-dependent dioxygenase enables efficient chemoenzymatic synthesis of manzacidin C and proline analogs, *J. Am. Chem. Soc.* 140 (3) (2018) 1165–1169.
- [104] C.R. Zwick, H. Renata, Harnessing the biocatalytic potential of iron-and α -ketoglutarate-dependent dioxygenases in natural product total synthesis, *Nat. Prod. Rep.* 37 (8) (2020) 1065–1079.
- [105] C.R. Zwick, M.B. Sosa, H. Renata, Characterization of a citrulline 4-hydroxylase from nonribosomal peptide GE81112 biosynthesis and engineering of its substrate specificity for the chemoenzymatic synthesis of enduracididine, *Angew. Chem. Int. Ed.* 58 (52) (2019) 18854–18858.